

## Ncr1-2A-Cre

Nomenclature C57BL/6Smoc-*Ncr1*<sup>em2(2A-iCre)Smoc</sup>

**Cat. NO.** NM-KI-190027

Strain State Repository Live

## **Gene Summary**

Gene Symbol Ncr1	Synonyms	Ly94; NKp46
	NCBI ID	<u>17086</u>
	MGI ID	1336212
	Ensembl ID	ENSMUSG00000062524
	Human Ortholog	NCR1

## **Model Description**

A 2A-iCre expression cassette was knocked into the Ncr1 gene stop codon site.

**Research Application**: Cre recombinase tool

\*Literature published using this strain should indicate: Ncr1-2A-Cre mice (Cat. NO. NM-KI-190027) were purchased from Shanghai Model Organisms Center, Inc..

## **Validation Data**






Fig1 Cre-mediated recombination in NK cells derived from spleen and peripheral blood of Ncr1-(2A-iCre);Rosa26 tdTomato mice.






Fig. 2: Detection of tdTomato (red) in some cells of the skin, kidney, submandibular gland(SMG) and brain drived from Ncr1icre/+; Rosa26tdTomato/+ mice.






Fig. 3 Detection of tdTomato (red) in various tissues of Ncr1icre/+; Rosa26tdTomato/+ mice. Cremediated recombination can be detected in some cells of skin, kidney, liver, thymus, spleen, submandibular gland(SMG), brain, pancreas, colon, intestine, brown fat, testis and epididymis. But tdTomato expression can not be observed in the lung, heart or muscle. (For more detailed information please contact our technical advisor.)